arXiv:1106.6311v2 [cond-mat.supr-con] 2 Dec 2011

Modulated superconductivity near Pomeranchuk instabilities in the spin channel

Daniel G. Barci[1] and Paulo S. A. Bonfim[1, 2]
[1]*Departamento de Física Teórica, Universidade do Estado do Rio de Janeiro, Rua São Francisco Xavier 524, 20550-013, Rio de Janeiro, RJ, Brazil.*
[2]*Universidade UnigranRio, Rua Prof. José de Souza Herdy, 1.160, 25071-202, Duque de Caxias,RJ, Brazil*

(Dated: December 5, 2011)

We study the competition between a Pomeranchuk instability in the spin channel with angular momentum $\ell = 1$ and an attractive interaction, favoring Cooper pair formation. We find, at mean-field approximation, that superconductivity strongly suppress the Pomeranchuk instability. Moreover, we have found a metastable modulated superconducting phase with similar characteristics of the FFLO state.

PACS numbers: 71.10.Hf,71.10.Ay,74.25.Dw,71.10.Pm

A Fermi liquid is, except in one dimension, a very stable state of matter[1]. At least two types of instabilities, related with attractive interactions, are known: Pomeranchuk[2] and superconducting instabilities. Pomeranchuk instabilities occur in the presence of two-body interactions containing a strong attractive component in the forward scattering channel with a definite angular momentum. In the context of Landau theory, the instability sets in when one or more dimensionless Landau parameters $F_\ell^{s,a}$, with angular momentum ℓ in the charge (s) or spin (a) channel, acquire large negative values.

Pomeranchuk instabilities in a charge sector spontaneously break rotational symmetry. In particular, an instability in the F_2^s channel produces an ellipsoidal deformation of the Fermi surface[3,4]. From a dynamical point of view, the resulting anisotropic ground state is a non-Fermi liquid, due to overdamped Goldstone modes that wipe out quasi-particle excitations[5–7].

In the spin channel, the ferromagnetic Stoner instability[8] occurs when F_0^a acquires large negative values, producing a divergence in the magnetic susceptibility. This phase transition preserves rotational symmetry, however it breaks time-reversal symmetry. Higher order angular momentum interactions produce anisotropic as well as isotropic phases. Several examples were studied in detail in refs. 9 and 10. The $\ell = 1$ channel have special interest. When $F_1^a < 0$, an ordered isotropic and time-reversal invariant phase is possible. This phase, called β-phase in ref. 9, dynamically generates a spin-orbit coupling. This is a very interesting possibility, since it allows the generation of spin-orbit couplings from many-body correlations, differently from the usual one-particle relativistic effect.

On the other hand, superconductivity is developed in the presence of a small attractive interaction in the particle-particle (BCS) channel. The superconducting (SC) state is generally characterized by a complex order parameter which breaks gauge symmetry, $\Delta_{\sigma,\sigma'}(\vec{r},\vec{r}') = \langle\psi^\dagger_\sigma(\vec{r})\psi^\dagger_{\sigma'}(\vec{r}')\rangle$, where the operator $\psi^\dagger_\sigma(\vec{r})$ creates an electron with spin σ at the position $\vec{r}$. The usual classification of $\Delta_{\sigma,\sigma'}(\vec{r},\vec{r}')$ as s-wave, d-wave, p-wave, etc, resides in the irreducible representations of the lattice point group. Also, the absence of spin-orbit interactions allows the additional differentiation between singlet and triplet order parameters. However, the SC state could break lattice translation and/or rotational symmetry. In that case, this classification is no more possible. One particular example is an oscillating order parameter like $\Delta_{\sigma,\sigma'}(\vec{r},0) = \Delta_{\sigma,\sigma'}\cos(\vec{q}\cdot\vec{r})$, where $\vec{q}$ is an ordering wave-vector. In a modulated superconducting state, proposed by Fulde, Ferrell, Larkin and Ovchinnikov[11,12] (FFLO state), the spatial modulation of the Cooper pairing is due to a mismatch of Fermi surfaces, produced by an external magnetic field (by Zeeman effect). FFLO states have also been proposed to occur in other scenarios like in imbalanced cold atoms with different species[13] and in heavy fermions systems when different orbitals hybridize under external pressure[14]. Also, for generally non-local BCS potentials, a modulated SC order parameter can coexist with charge density waves[15]. Recently, a striped order parameter, called "Pair Density Wave, (PDW)", was proposed[16–18] to explain anomalous transport properties observed in cuprates superconductors.

Although Pomeranchuk and BCS instabilities are generated by attractive interactions, they are competing ones. A superconducting gap suppresses Fermi surface deformation. A detailed example of this effect was studied in refs. 19 and 20, where the competition between d-wave Pomeranchuk instabilities and d-wave superconducting order parameter was considered.

In the spin channel, a stronger competition is expected, since, in general, magnetic order is expelled from a superconductor state. However, it is possible to have instabilities with higher angular momentum that preserve time-reversal invariance. In this paper, we analyze an example of this class of systems. In particular, the β-phase introduced in ref. 9, opens the possibility of the formation of Cooper pairs with zero helicity and finite momentum, possibly producing a modulated superconducting state.

The main point of this paper is to report on the com-



FIG. 1. Phase diagram. $\bar{\Delta}_0$ is the dimensionless isotropic BCS gap $\bar{\Delta}_0 = b\Delta_0/v_F k_F$. $F_1^a = f_1^a N(0)$ is the dimensionless antisymmetric Landau parameter with angular momentum $\ell = 1$. The solid line, separating the β and the SC phase, is a first order line. The dashed line indicate the onset of a metastable modulated SC phase.

peting character of Cooper pairing and a Pomeranchuk instability in the $\ell = 1$ spin channel. Our model is based on a 2D Fermi liquid with interactions in the F_1^a channel and a general attracting interaction favoring Cooper pairing. The main results are summarized in the phase diagram of fig. (1), obtained in a self-consistent mean-field approximation described below. In the figure, $\Delta_0 \sim e^{-1/g}$ is a parametrization of the BCS coupling constant g. Physically, it represents the gap of the uniform s-wave superconductor in the absence of any Pomeranchuk instability. For $F_1^a > -2$, the system is in the usual s-wave SC state. When $F_1^a < -2$ the Pomeranchuk instability sets in and the mean-field ground state at low Δ_0 is the so called β-phase consisting on a Fermi liquid with an induced spin-orbit interaction. For higher values of Δ_0, s-wave superconductivity strongly suppresses the β-phase. They are separated by a coexisting line, signaling a first order phase transition. Moreover, for greater values of the BCS coupling, a metastable modulated superconducting phase appears with a finite wave-vector $\vec{q}$, given roughly by $|\vec{q}| \sim k_F\sqrt{1-2/|F_1^a|}$.

In the rest of the paper we present our model and sketch the mean-field calculation leading to the phase diagram of fig. (1). A spin-current operator with orbital angular momentum $\ell = 1$ is given by

$$V^{a\mu}(x) = -i\psi^\dagger(x)\sigma^\mu\hat{\nabla}_a\psi(x) \ , \qquad (1)$$

where ψ is a two-component spinor, σ^μ are the Pauli matrices, and $a = x, y$. $\hat{\nabla}e^{ikx} = i\frac{\vec{k}}{|\vec{k}|}e^{ikx}$.

Following ref. 9, we write a Hamiltonian with forward two-body interactions in this channel

$$H = \int d^2x\, \psi^\dagger(x)\left(\epsilon(-i\nabla) - \mu\right)\psi(x) + \\ + \frac{1}{2}\int d^2x d^2x'\, f_1^a(x-x')V^{a\mu}(x)V^{a\mu}(x') \ . \qquad (2)$$

The Fourier transform of $f_1^a(x)$ is given by $f_1^a(k) = f_1^a/(1+\kappa|f_1^a|k^2)$, defining, in this way, an effective interaction range $r = \sqrt{\kappa|f_1^a|}$. For a particle-hole symmetric system we consider the following expansion of the dispersion relation around a circular Fermi surface,

$$\epsilon(k) - \mu = \vec{v}_F\cdot[\vec{k}-\vec{k}_F] + \frac{b}{(v_F k_F)^2}\left(\vec{v}_F\cdot[\vec{k}-\vec{k}_F]\right)^3 \ , \qquad (3)$$

where the dimensionless parameter b measures the effective curvature of the band near the Fermi surface and we have ignored terms proportional to $(k-k_F)^5$.

The β-phase is defined by the mean-field Hamiltonian

$$H_{MF} = \int \frac{d^2k}{(2\pi)^2}\, \psi^\dagger(k)\left(\epsilon(k) - \mu - \bar{n}\vec{\sigma}\cdot\hat{k}\right)\psi(k), \qquad (4)$$

where $\bar{n}$ is determined self-consistently by

$$\bar{n} = -\frac{1}{2}f_1^a(0)\int \frac{d^2k}{(2\pi)^2}\langle\psi^\dagger(k)(\vec{\sigma}\cdot\hat{k})\psi(k)\rangle \ . \qquad (5)$$

This mean-field theory is valid when $k_F\sqrt{\kappa|f_1^a|} \gg 1$, *i. e.* , when the range of the interaction is much larger than the interparticle distance.

In references 9 and 10, eq. (5) was solved by considering a Fermi liquid ground state. When $\bar{n} \neq 0$ the spectrum of H_{MF} splits into two opposite chiralities with dispersions $\epsilon^\uparrow = \epsilon(k) - (\mu + \bar{n})$ and $\epsilon^\downarrow = \epsilon(k) - (\mu - \bar{n})$, as shown in figure (2). The Fermi momentum of each branch is given by $k_F^{\uparrow\downarrow} = k_f \pm q/2$ and the relation between q and $\bar{n}$ is computed using eq. (3),

$$\frac{v_F q}{2} = \bar{n} - \frac{b}{(v_F k_F)^2}\,\bar{n}^3 + O(\bar{n}^5) \ , \qquad (6)$$

(we have renormalized the chemical potential to keep the density constant).

The ordered phase is characterized by a spontaneously generated spin-orbit coupling with the global rotation invariance unbroken. The spin and orbital angular momentum are not conserved independently, however the total angular momentum $\vec{J} = \vec{L} + \vec{S}$ is conserved. In the helicity base $\zeta = (\zeta_\uparrow, \zeta_\downarrow)$, where the operator $\hat{k}\cdot\vec{\sigma}$ is diagonalized with eigenvalues ± 1, the mean-field Hamiltonian takes the simpler form

$$H_{MF} = \int \frac{d^2k}{(2\pi)^2}\, \zeta^\dagger(k)\left(\epsilon(k) - \mu - \bar{n}\sigma^z\right)\zeta(k) \ , \qquad (7)$$

with the self-consistent equation

$$\bar{n} = -\frac{1}{2}f_1^a(0)\int \frac{d^2k}{(2\pi)^2}\langle\zeta^\dagger\sigma^z\zeta\rangle \ . \qquad (8)$$



$\zeta^\dagger_{\uparrow,k+q/2}$
$\vec{k}^\downarrow_F$
$\vec{k}_F$
q
$\vec{k}^\uparrow_F$
$\zeta^\dagger_{\downarrow,-k+q/2}$

FIG. 2. Fermi Surface splitting in the helicity base and pairing in the β-phase.

A BCS type interaction in the $\psi, \psi^\dagger$ base induces different types of interactions in the β-phase, where the usual Fermi liquid properties are lost. In the helicity base $\zeta, \zeta^\dagger$, intra-band p-wave pairing, as well as inter-band finite momentum pairing, are induced. These interactions are competing with each other and with interactions in the forward scattering (Pomeranchuk) channel. In this work, we study the competition between an inter-band SC order parameter and the β-instability, leaving a complete study of the phase diagram, including other Cooper pair symmetries and fluctuations, for a future presentation.

We consider a simple model where we introduce the following attractive interaction favoring Cooper pair formation with linear momentum $\vec{q}$,

$$H_{SC} = \int \frac{d^2k}{(2\pi)^2} \Delta_q \, \zeta^\dagger_{\uparrow,k+q/2} \zeta^\dagger_{\downarrow,-k+q/2} + \text{h.c.} \,, \quad (9)$$

Note that $\vec{q}$ (eq. (6)) is given by the Pomeranchuk instability and should be determined self-consistently. The "gap equation" that complements eq. (9) is

$$\Delta_q = -g \int \frac{d^2k}{(2\pi)^2} \langle \zeta_{\downarrow,-k+q/2} \zeta_{\uparrow,k+q/2} \rangle \,. \quad (10)$$

To find a self-consistent solution, it is necessary to solve eqs. (8), (10) and (6) simultaneously, where the expectation values are taken with the Hamiltonian $H = H_{MF} + H_{SC}$.

Diagonalizing H, considering $f_1^a < 0$, we find at zero temperature

$$\bar{n} = \frac{|f_1^a|}{4} \int \frac{d^2k}{(2\pi)^2} \left\{ \Theta(-\omega_+) - \Theta(\omega_-) \right\} \,, \quad (11)$$

$$\Delta_q = -g \int \frac{d^2k}{(2\pi)^2} \frac{\Delta_q}{2\xi} \left\{ \Theta(-\omega_+) - \Theta(-\omega_-) \right\} \,, \quad (12)$$

where Θ is the usual Heaviside function and single-particle excitations read

$$\omega_\pm = \left(\frac{\epsilon^\uparrow_{k+q/2} - \epsilon^\downarrow_{k-q/2}}{2} \right) \pm \xi \,, \quad (13)$$

with

$$\xi = \sqrt{ \left(\frac{\epsilon^\uparrow_{k+q/2} + \epsilon^\downarrow_{k-q/2}}{2} \right)^2 + |\Delta_q|^2 } \,. \quad (14)$$

It is simple to realize that the s-wave BCS order parameter $\bar{n} = 0, \Delta_q = \Delta_0$ is always a solution of eqs. (11) and (12). On the other hand, in the absence of superconductivity, $\Delta_0 = 0$, a solution with finite $\bar{n} \sim q$ has lower energy for $F_1^a < -2$, leading to the β-phase. In this regime and for small values of the BCS coupling constant, the β-phase has always smaller energy than the uniform superconducting one. For stronger superconducting coupling, the β-phase is suppressed and the mean-field ground state is a uniform superconducting phase, as shown in figure (1). The first order line that separates these two phases was evaluated by equating the mean-field Hamiltonians, $\langle H \rangle_{\Delta_0} = \langle H \rangle_\beta$, computed in the two possible solutions: uniform superconductor $\{\bar{n} = 0, \Delta_q = \Delta_0\}$ and the β-phase $\{\bar{n} \sim (v_F k_F / b^{1/2}) \sqrt{1 - 2/|F_1^a|}, \Delta_q = 0\}$, respectively. Note that the band curvature at the Fermi surface (b) is essential to stabilize the β-phase.

Now, we explore the interesting possibility of a modulated solution $\bar{n} \sim q \neq 0, \Delta_q \neq 0$. The $\vec{k}$-integrals in equations (11) and (12) are strongly constrained by the Heaviside functions. The main contribution to eq. (11) comes from the region $\omega_+ < 0$. Written in polar coordinates (k, θ), with $\cos\theta = (\vec{k} \cdot \vec{q})/kq$, and first integrating over k, we get

$$\bar{n} = |F_1^a| \bar{n} \int_{\theta^-}^{\theta^+} \frac{d\theta}{2\pi} \times$$
$$\times \sqrt{ \left(1 - \frac{b}{(v_F k_F)^2} \bar{n}^2 \cos\theta \right)^2 (1 + \cos\theta)^2 - x^2 } \quad (15)$$

where we have defined the dimensionless quantity $x = \Delta_q / \bar{n}$. $\theta^\pm(x, \bar{n})$ are the integration limits which keep the argument of the square root positive. Near the critical point, we can expand the integrand in powers of $\bar{n}/v_F k_F << 1$, obtaining

$$\bar{n} = \frac{|F_1^a|}{2} \bar{n} \left[C_0(x) - \frac{b}{(v_F k_F)^2} \bar{n}^2 C_1(x) \right] + O\left(\left(\frac{\bar{n}}{v_F k_F} \right)^5 \right), \quad (16)$$

with $C_0(x) \sim 1 - (x/2)^{3/2}$, $C_1(x) = 1 + x \ln(1 + x/5)$ and $F_1^a = f_1^a N(0)$, where $N(0)$ is the density of states at the Fermi surface.

On the other hand, the integral in eq. (12) has the usual ultraviolet divergence of the BCS gap equation. A convenient way to deal with this integral is to sum and subtract $\Theta(\omega_-)$ in the second term and then to subtract the identity

$$-1 + g \int \frac{d^2k}{(2\pi)^2} \frac{1}{2\xi^0} \equiv 0 \quad (17)$$

from the first term of eq. (12). $\xi^0 = \sqrt{\epsilon_k^2 + |\Delta_0|^2}$ and Δ_0 is the uniform superconducting gap in the absence of

Fermi surface splitting. With this, eq. (12) is rewritten as

$$\frac{N(0)}{2}\ln\left|\frac{\Delta_0}{\Delta_q}\right| = \int \frac{d^2k}{(2\pi)^2}\frac{1}{2\xi}\left\{\Theta(-\omega_+)+\Theta(\omega_-)\right\} \ . \quad (18)$$

In this way, the above integral is controlled. The coupling constant g and the ultraviolet cut-off are both contained in the definition of Δ_0 (eq. (17)). As before, near the Fermi surface, where interactions are important, $\Theta(\omega_-)=0$. Performing the remaining integral over k, the gap equation is written as

$$\ln\left|\frac{\Delta_0}{\Delta_q}\right| = \Gamma(x) \ , \quad (19)$$

where $x=\Delta_q/\bar{n}$, and we have defined the function

$$\Gamma(x) = 2\int_{-\bar{\theta}(x)}^{\bar{\theta}(x)}\frac{d\theta}{2\pi}\,\sinh^{-1}\left(\frac{\sqrt{(1+\cos\theta)^2-x^2}}{x^2}\right) \ , \quad (20)$$

with $\bar{\theta}(x)=\cos^{-1}(x-1)$ for $x\leq 2$ and $\Gamma(x>2)=0$. Therefore, to look for modulated superconductivity we need to solve eqs. (16) and (19) self-consistently. Firstly, we note that for $x>2$ the only solution, $\{\bar{n}=0,\Delta_q=\Delta_0\}$, corresponds to the uniform superconducting phase. This imposes a lower limit for modulated superconductivity since, for obtaining a non-trivial solution, we must have $\Delta_q<2\bar{n}<\Delta_0$. However, for any value of the parameter $\Delta_0>2\bar{n}$, the uniform superconducting mean-field energy is always lower than the modulated one, $\langle H\rangle_{\Delta_0}<\langle H\rangle_{\Delta_q}$. On the other hand, there is a region of the phase diagram, shown over the dashed line in figure (1), in which $\langle H\rangle_{\Delta_q}<\langle H\rangle_{\beta}$. Therefore, modulated superconductivity appears as a metastable phase for finite values of Δ_0 and $|F_1^a|>2$. The precise location of the onset of metastability depends on the relation between the band-width, in which interactions are relevant (the energy cut-off), and the band-curvature at the Fermi surface.

Summarizing, we have studied, in the context of Fermi liquids, the competition between a Pomeranchuk instability in the spin channel with angular momentum $\ell=1$, triggered by the Landau Parameter F_1^a and an attractive interaction favoring Cooper pair formation. We built up a zero temperature mean-field phase diagram in terms of the two parameters of the model: F_1^a and g, parametrized by Δ_0. We found that, as observed in other channels of Pomeranchuk instabilities[19,20], the superconducting gap reduces the tendency to Fermi surface deformation. Moreover, we have found a metastable modulated superconductor, enhanced for larger values of $|F_1^a|$.

The phase diagram of figure (1) could be modified by several reasons. On the one hand, we have ignored intra-band homogeneous pairing that surely competes with the inter-band pairing considered in this paper. On the other hand, long wave-length fluctuations will certainly modify mean-field results. There are two types of important fluctuations to be considered. In the β-phase, there are three branches of Goldstone modes transforming with $SO(3)$ group, associated with the breaking of relative spin-orbit symmetry[10]. The longitudinal mode has a linear dispersion relation while the transverse ones are responsible for a Lifshitz instability, signaling a tendency to a modulated ground-state with parity symmetry breaking. This effect, in the presence of pairing interactions, should probably enhance the modulated SC order.

D.G.B. is in debt with Eduardo Fradkin for very useful discussions. The Brazilian agencies *“Conselho Nacional de Desenvolvimento Científico e Tecnológico”, (CNPq)* and the *“Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro” , (FAPERJ)*, are acknowledged for partial financial support.

---

[1] P. Nozieres and D. Pines, *The Theory of Quantum Liquids* (Perseus Books, 1999).

[2] I. I. Pomeranchuk, Sov. Phys. JETP **8**, 361 (1958).

[3] V. Oganesyan, S. A. Kivelson, and E. Fradkin, Phys. Rev. B **64**, 195109 (2001).

[4] D. G. Barci and L. E. Oxman, Phys. Rev. B **67**, 205108 (2003).

[5] M. J. Lawler, D. G. Barci, V. Fernández, E. Fradkin, and L. Oxman, Phys. Rev. B **73**, 085101 (2006).

[6] M. A. Metlitski and S. Sachdev, Phys. Rev. B **82**, 075127 (2010).

[7] D. G. Barci, M. Trobo, V. Fernández, and L. E. Oxman, Phys. Rev. B **78**, 035114 (2008).

[8] Y. Zhang and S. Das Sarma, Phys. Rev. B **72**, 115317 (2005).

[9] C. Wu and S.-C. Zhang, Phys. Rev. Lett. **93**, 036403 (2004).

[10] C. J. Wu, K. Sun, E. Fradkin, and S.-C. Zhang, Phys. Rev. B **75**, 115103 (2007).

[11] P. Fulde and R. A. Ferrell, Phys. Rev. **135**, A550 (1964).

[12] A. I. Larkin and Y. N. Ovchinnikov, Zh. Eksp. Teor. Fiz. **47**, 1136 (1964), (Sov. Phys. JETP. **20**, 762 (1965)).

[13] L. Radzihovsky and A. Vishwanath, Phys. Rev. Lett. **103**, 010404 (2009).

[14] I. T. Padilha and M. A. Continentino, Journal of Physics: Condensed Matter **21**, 095603 (2009).

[15] F. Loder, A. P. Kampf, and T. Kopp, Phys. Rev. B **81**, 020511 (2010).

[16] E. Berg, E. Fradkin, and S. A. Kivelson, Nature Physics **5**, 830 (2009).

[17] E. Berg, E. Fradkin, S. A. Kivelson, and J. M. Tranquada, New J. Phys. **11**, 115004 (2009).

[18] D. G. Barci and E. Fradkin, Phys. Rev. B **83**, 100509 (2011).

[19] C. J. Halboth and W. Metzner, Phys. Rev. Lett. **85**, 5162 (2000).

[20] H. Yamase and W. Metzner, Phys. Rev. B **75**, 155117 (2007).